

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS✓
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Branched maltodextrins and process for their preparation

No. Publication (Sec.) : EP1006128

Date de publication : 2000-06-07

Inventeur : FOUACHE CATHERINE (FR); DUFLOT PIERRICK (FR); LOOTEN PHILIPPE (FR)

Déposant : ROQUETTE FRERES (FR)

Numéro original : ☐ EP1006128

No. d'enregistrement : EP19990403024 19991203

No. de priorité : FR19980015344 19981204

Classification IPC : C08B30/18

Classification EC : C08B30/18, A23G3/00K, A23K1/16L, A23L1/09D, A23L1/09E

Brevets

correspondants : AU6303099, ☐ FR2786775, ☐ JP2000169502

Cited patent(s): US4840807; JP48067447; JP7252301; JP51088645; JP3137102

Abrégé

Branched maltodextrins (I) having 22-35% of 1-6 glucoside linkages, a reducing sugar content of less than 20%, a polymolecularity index (Mw/Mn) of less than 5 and a number-average molecular weight (Mn) of up to 4500 are new. Independent claims are also included for: (1) an acariogenic composition comprising (I) and at least one polyol; (2) a process for preparing (I), comprising: (a) preparing an acidified starch with a moisture content below 5%; (b) heating the starch in a continuous thin-film reactor at 120-300 degrees C; (c) recovering, purifying and preferably concentrating the resulting branched starch derivatives; and (d) fractionating the starch derivatives according to Mn.

Données extraites de la base de données esp@cenet - I2